

1st Semester
1439 – 1440 | 2018 – 2019

ALWAYS LEARNING

P Pearson

Taibah University
The Unified Scientific Track

Lecture Presentation

Chapter 7

The Chemistry of Life: Organic and Biological Chemistry

Topic 20

- Introduction to Organic Chemistry
- Hydrocarbons
- Alkanes & Cycloalkanes

Introduction To Organic Chemistry

- Organic chemistry is an old interesting branch of chemistry, but has only started in the 19th century as a science in its modern sens.
- **Organic Chemistry** is the chemistry of **carbon element**. Carbon forms strong chemical bonds to other carbon atoms and to many other elements.
- Because of its versatility in forming covalent bonds, millions of carbon compounds are known.
- The existence of a great number of different organic compounds has raised up the need to classify them into “**families**”.
- Carbon always forms **four covalent bonds** (four shared pairs of electrons) that may be present as:
 - 4 single bonds per atom,
 - 1 single and 1 triple bond, or
 - 2 single and 1 double bond,
 - 2 double bonds.

Hydrocarbons

- The family of “**Hydrocarbons**” is the simplest family of organic compounds, containing only hydrogen and carbon atoms.
- **Hydrocarbons** are **non-polar** molecules, **insoluble** in water and soluble in non-polar solvents.
- Hydrocarbons have low melting and boiling points.
- There are four basic types of hydrocarbons:
 - **Alkanes** (C–C)
 - **Alkenes** (C=C)
 - **Alkynes** (C≡C)
 - **Aromatic hydrocarbons** ()

Hydrocarbons (contain only carbon and hydrogen)

Alkanes
(only C—C bonds)

Alkenes
(C=C bond)

Alkynes
(C≡C bond)


Aromatic
(contains benzene ring)

Aliphatic

Generic Formula*


C_nH_{2n+2}

Example




Generic Formula*

C_nH_{2n}




Generic Formula*

C_nH_{2n-2}



Benzene Formula

C_6H_6





Benzene

Alkanes (C—C)

Alkane



Ethane



The general formula of **alkanes** is **$\text{C}_n\text{H}_{2n+2}$**

Alkanes are known as “**saturated hydrocarbons**” that contain only single bonds (C—C).

- Every carbon atom has **4 single bonds (C—C)**, no double nor triple bonds are present in alkanes.




Representing Bonding Connections

There are 3 ways to represent bonding connections:

Example: The different bonding connections for n-butane ($\mathbf{C_4H_{10}}$):

1. Expanded Structure:



2. Condensed Structure:



3. Stick (Carbon Skeleton):



Properties of Alkanes

- Boiling points of **Alkanes** increase as chain length increases:

TABLE 24.2 • First Ten Members of the Straight-Chain Alkane Series






Molecular Formula	Condensed Structural Formula	Name	Boiling Point (°C)
CH ₄	CH ₄	Methane	−161
C ₂ H ₆	CH ₃ CH ₃	Ethane	−89
C ₃ H ₈	CH ₃ CH ₂ CH ₃	Propane	−44
C ₄ H ₁₀	CH ₃ CH ₂ CH ₂ CH ₃	Butane	−0.5
C ₅ H ₁₂	CH ₃ CH ₂ CH ₂ CH ₂ CH ₃	Pentane	36
C ₆ H ₁₄	CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	Hexane	68
C ₇ H ₁₆	CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	Heptane	98
C ₈ H ₁₈	CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	Octane	125
C ₉ H ₂₀	CH ₃ CH ₂ CH ₃	Nonane	151
C ₁₀ H ₂₂	CH ₃ CH ₂ CH ₃	Decane	174

© 2012 Pearson Education, Inc.


Isomers of Alkanes

➤ **Isomers**: compounds that have the same molecular formula but **different chemical structures** (i.e. different order of bonding).

TABLE 24.3 • Isomers of C_4H_{10} and C_5H_{12}

Systematic Name (Common Name)	Structural Formula	Condensed Structural Formula	Space-filling Model	Melting Point (°C)	Boiling Point (°C)
Butane (<i>n</i> -butane)		$CH_3CH_2CH_2CH_3$		-138 °C	-0.5 °C
2-Methylpropane (isobutane)		$CH_3-CH(CH_3)_2$		-159 °C	-12 °C
Pentane (<i>n</i> -pentane)		$CH_3CH_2CH_2CH_2CH_3$		-130 °C	+36 °C
2-Methylbutane (isopentane)		$CH_3-CH(CH_3)-CH_2-CH_3$		-160 °C	+28 °C
2,2-Dimethylpropane (neopentane)		$CH_3-C(CH_3)_3$		-16 °C	+9 °C

Nomenclature of Organic Compounds



➤ The names of organic compounds contain three parts:

- **Prefix:** This tells what substituent groups are attached to the chain.
- **Base:** This tells how many carbons are there in the longest continuous carbon chain.
- **Suffix:** This tells what type of compound it is (the family)

Nomenclature of Alkanes


➤ The Base Names:

Alkane Nomenclature

Number of carbon atoms	Base Name	Alkane Formula	Name of alkane	Name of alkyl group (R)	Alkyl (R) Formula
1	meth-	CH_4	methane	methyl	CH_3-
2	eth-	CH_3CH_3	ethane	ethyl	CH_3CH_2-
3	prop-	$\text{CH}_3\text{CH}_2\text{CH}_3$	propane	propyl	$\text{CH}_3\text{CH}_2\text{CH}_2-$
4	but-	$\text{CH}_3(\text{CH}_2)_2\text{CH}_3$	butane	butyl	$\text{CH}_3(\text{CH}_2)_2\text{CH}_2-$
5	pent-	$\text{CH}_3(\text{CH}_2)_3\text{CH}_3$	pentane	pentyl	$\text{CH}_3(\text{CH}_2)_3\text{CH}_2-$
6	hex-	$\text{CH}_3(\text{CH}_2)_4\text{CH}_3$	hexane	hexyl	$\text{CH}_3(\text{CH}_2)_4\text{CH}_2-$
7	hept-	$\text{CH}_3(\text{CH}_2)_5\text{CH}_3$	heptane	heptyl	$\text{CH}_3(\text{CH}_2)_5\text{CH}_2-$
8	oct-	$\text{CH}_3(\text{CH}_2)_6\text{CH}_3$	octane	octyl	$\text{CH}_3(\text{CH}_2)_6\text{CH}_2-$
9	non-	$\text{CH}_3(\text{CH}_2)_7\text{CH}_3$	nonane	nonyl	$\text{CH}_3(\text{CH}_2)_7\text{CH}_2-$
10	dec-	$\text{CH}_3(\text{CH}_2)_8\text{CH}_3$	decane	decyl	$\text{CH}_3(\text{CH}_2)_8\text{CH}_2-$

Nomenclature of Alkanes

1. Find the **longest continuous chain** of carbon atoms in the molecule and use this chain as the **base name** (see the table of base names).





2. Number the carbon atoms in the longest chain, beginning with the end **nearest to a substituent**.
3. Name each substituent (**prefixes**)
4. Begin the name with the number or numbers of carbon atoms to which each substituent is bonded.
5. When two or more substituents are present, list them **alphabetically**

Names of Substituent Groups (Branches)

➤ Carbon Groups (alkyl groups, R):


- Methyl CH_3-
- Ethyl CH_3CH_2-
- Propyl $\text{CH}_3\text{CH}_2\text{CH}_2-$
- Butyl $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2-$



➤ Halogens:

- Fluoro $\text{F}-$
- Chloro $\text{Cl}-$
- Bromo $\text{Br}-$
- Iodo $\text{I}-$





Names of Substituent Groups (Branches)




3-Ethyl-2,4,5-trimethylheptane

If there is more than one type of substituent in the molecule, list them “**alphabetically**”.

Nomenclature of Alkanes: Exercises




✓ 4-Ethyl-3-methylocatane
✗ NOT 3,3-Methyl-4-ethyloctane




✓ 3,5,7-Trimethyldecane
✗ NOT 4,6,8-Trimethyldecane


Nomenclature of Alkanes: Exercises




Chloromethane




Bromoethane




1-Iodopropane




2-Iodopropane




1-Bromobutane




2-Bromobutane



1-Bromo-2methylpropane



1-chlorohexane










1-Chloro-2-methylbutane

Cycloalkanes



- Carbon can also form **cyclic (ringed)** structures.

The general formula of **cycloalkanes** is C_nH_{2n}



- Six-membered rings are the most stable cyclic compounds.



Nomenclature of Cycloalkanes




- ✓ 1,3-dimethylcyclohexane
- ✗ (not 1,5-dimethylcyclohexane)



Earlier letter -----> lower number

- ethyl group at C1
- methyl group at C3



- ✓ 1-ethyl-3-methylcyclohexane
- ✗ (not 3-ethyl-1-methylcyclohexane)




Reactions of Alkanes

- Alkanes are mainly used as **non-polar solvents**.
- Most alkanes are relatively **unreactive** at room temperature, because they contain only C-C and C-H bonds.
- However, alkanes are not completely inert. One of their important reactions is their **combustion** in oxygen, making them important **fuels** and a source of thermal energy:
- **Example:** the combustion of ethane:



Assessment

1. Give the name or structural formula, as appropriate:



(d) 2-methylheptane

(e) 2,2-dimethylpentane







(f) 4-ethyl-2,3-dimethyloctane


(g) 4-ethyl-1,1-dimethylcyclohexane

(h) 1,2-dimethylcyclohexane

(i) $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{C}(\text{CH}_3)_3$

2. Which of the following pairs of compounds are isomers?





Chapter 7

The Chemistry of Life: Organic and Biological Chemistry

Topic 21

- Alkenes
- Alkynes
- Aromatic Hydrocarbons





Taibah University
The Unified Scientific Track

ALWAYS LEARNING

P Pearson

Alkenes (C=C)

Alkene Ethylene $\text{CH}_2=\text{CH}_2$




The general formula of **alkenes** is C_nH_{2n}

Alkenes are **unsaturated hydrocarbons** that contain at least one double bond (**C=C**).


- The simplest alkene is $\text{CH}_2=\text{CH}_2$, called [ethene](#) (IUPAC) or [ethylene](#) (common name).

Alkene



Ethylene: C_2H_4
(fewer hydrogens—*unsaturated*)


Alkane



Ethane: C_2H_6
(more hydrogens—*saturated*)

Structure of Alkenes: *cis/trans* Isomers

- Unlike alk^{anes}, alk^{enes} cannot rotate around the C=C bond:





cis/trans geometrical isomerism:

- ***cis*-Alkenes**: have the R groups on the same side of the double bond plan.

- ***trans*-Alkenes**: have the R groups on opposite sides of the double bond plan.


- Geometric isomers can differ significantly from each other in chemical behaviour.




Nomenclature of Alkenes

Example

Give the IUPAC name of the following alkene:



Step [1] Find the longest chain that contains *both* carbon atoms of the double bond.




hexane \dashrightarrow hexene

- Change the *-ane* ending of the parent alkane to *-ene*.


Step [2] Number the carbon chain to give the double bond the lower number, and apply all other rules of nomenclature.

[a] Number the chain, and name using the *first number* assigned to the C=C.



2-hexene

[b] Name and *number* the substituents.





three methyl groups at C2, C3, and C5


Answer: 2,3,5-trimethyl-2-hexene

Nomenclature of Alkenes

Start numbering here.




Answer: 3-methyl-1-butene



Both C's of the C=C are NOT contained in this long chain.

✗ *Incorrect*







7 C's → heptene

Both C's of the C=C are contained in this long chain.



✓ *Correct: 2-ethyl-1-heptene*

Nomenclature of Alkenes




Note: If an alkene contains **two or more** double bonds, the location of each is indicated by numerical prefix, and the ending of the name is altered to identify the number of double bonds: diene(two), triene (three):

Example: CH2=CH-CH2-CH=CH2 is named: **1,4-penta^{diene}**.

Addition Reactions of Alkenes


- One important reaction of alkenes is the **Addition Reaction**:
 - In which, two atoms (e.g., bromine) add across the double bond.
 - One π -bond (from C=C) and one σ -bond (from Br–Br) are replaced by two σ -bonds (2 C–Br); therefore, ΔH is negative.



© 2012 Pearson Education, Inc.

Alkynes (C≡C)



Alkyne Acetylene CH≡CH




The general formula of **alkynes** is **C_nH_{2n-2}**

Alkynes are **unsaturated hydrocarbons** that contain at least one triple bond (**C≡C**).

- The simplest alkyne is H-C≡C-H, called [ethyne](#) (IUPAC) or [acetylene](#) (common name).




terminal alkyne



internal alkyne


Nomenclature of Alkynes



4-methyl-2-pentyne



- The method for naming alkynes is **similar** to that for naming alkenes.
- However, the suffix *-yne* is used rather than *-ene*.

Nomenclature of Alkynes



Addition Reactions of Alkynes

- Alkynes undergo many of the same reactions that alkenes do.
- As with alkenes, the drive for the addition reaction is the replacement of π -bonds by σ -bonds.



Aromatic Hydrocarbons

- **Aromatic Compound:** A hydrocarbon that contains one or more benzene-like rings.
- **Benzene** (C_6H_6) is the simplest and the most important aromatic hydrocarbon.
- It contains three alternated single/double bonds.
- Compared to alkenes, benzene is very stable and unreactive towards normal reagents.



Benzene



Naphthalene



Anthracene




Pyrene

Some Common Aromatic Hydrocarbons


- Some important mono-substituted benzene compounds have **common names** that you must learn:

- Structure:




- Common Name:

toluene



phenol



aniline

- IUPAC Name:

(methylbenzene)

(hydroxybenzene)

(aminobenzene)

*Students shall carefully memorize these examples.

Substitution Reactions of Benzene

- It reacts differently to alk^{enes}, yielding **substitution** products instead of **addition** ones.
- Benzene reacts **slowly** with Br_2 , producing the bromobenzene as a **substitution** product.
- **Addition** products are **NOT** formed.



Assessment

1. Name or write the condensed structural formula for the following compounds:

a) *trans*-2-pentene




b) 2,5-dimethyl-4-octene

c) 1,1-dichloro-1-butene

d) 1,4-dichlorobenzene

e) 2,4-dichloro-2-butene

f) 4,5-dimethyl-2-pentyne








2. Identify the type of the following hydrocarbons (alkane, alkene, or alkyne)

a) C₄H₈ b) C₄H₆ c) C₅H₁₂ d) C₇H₁₄

e) C₈H₁₆ f) C₁₈H₃₈ g) C₆H₁₀ h) C₁₀H₂₂

3. In the following carbon skeletons, how many hydrogen atoms shall be bonded to the carbon marked with a *?





1st Semester
1439 – 1440 | 2018 – 2019



Taibah University
The Unified Scientific Track

ALWAYS LEARNING

P Pearson

Chapter 7

The Chemistry of Life: Organic and Biological Chemistry

Topic 22

Organic Functional Groups:

Alcohols, Ethers, Aldehydes,

Ketones, Carboxylic Acids,

Esters, Amines & Amides

7.3 Organic Functional Groups


- The **Functional Group** is the active part of an organic molecule, where reactions tend to occur.

For example, the double bond C=C is the functional group of alkenes and the triple bond C≡C is the functional group of alkynes.

- When drawing the structure of some organic molecule, the **alkyl** parts are represented by **R** (**R** = CH_3- , CH_3CH_2- , $\text{CH}_3\text{CH}_2\text{CH}_2-$...) which are unreactive, giving rise to the functional groups to react.
 - ✓ When the present alkyl groups are different, they are represented as R, R', R" or R¹, R², R³,


Alcohols (R-OH)

- **Alcohols** are organic compounds, containing one or more (**-OH**) groups (called either the hydroxyl group or the alcohol group).
- The systematic names for alcohols ends with **-ol**.
- The O-H bond is polar, so alcohols are more soluble in polar solvents than are hydrocarbons.




Structure, Classes and Properties of Alcohols


- **Alcohols** are classified according to the number of carbon atoms bonded to the “C” carrying the “OH” group:



A **primary** (1°) alcohol has an OH group on a C which is bonded to another **1 C atom + 2 H atoms**.




A **secondary** (2°) alcohol has an OH group on a C which is bonded to other **2 C atoms + 1 H atoms**.




A **tertiary** (3°) alcohol has an OH group on a C which is bonded to other **3 C atoms + no H atoms**.

Naming of Alcohols


- Alcohols are named from the hydrocarbon parent; The suffix is changed to ***-ol*** and a number designates the carbon to which the hydroxyl is attached.



2-Propanol
Isopropyl alcohol;
rubbing alcohol




2-Methyl-2-propanol
t-Butyl alcohol




1,2-Ethanediol
Ethylene glycol



Phenol




1,2,3-Propanetriol
Glycerol; glycerin




Cholesterol

Naming of Alcohols

Exercise: Give the IUPAC name for the following compound:




✓ **Step 1:** Name the longest carbon chain attached to the **—OH** group by replacing the “**e**” in the corresponding alk**ane** name with **ol**.




Naming of Alcohols

✓ **STEP 2:** Number the chain starting at the end nearer to the **—OH** group.




✓ **STEP 3:** Give the location and name for each substituent relative to the **—OH** group.




Oxidation Reactions of Alcohol

- The partial oxidation of **primary alcohols** produce the corresponding **aldehydes**, while the further oxidation produces **carboxylic acids**.
- The partial oxidation of **secondary alcohols** produce the corresponding **ketones**.
- The oxidation of a tertiary alcohol is **not possible**.




Ethers (R–O–R')

Ethers: compounds in which two hydrocarbon groups (**R**) are bonded to one oxygen atom.




- Ethers can be formed from two molecules of alcohol by splitting out a molecule of water (**Condensation Reaction**). This reaction is catalyzed by sulfuric acid.



Ethers tend to be quite **unreactive**. Therefore, they are common solvents for organic reactions.






Diethyl ether



Tetrahydrofuran (THF)


Some **Ethers** are used as medical “**anesthetics**” that inhibit pain signals to the brain during surgeries.

Aldehydes (R-CHO) and Ketones (R-CO-R')




R & R'
Alkyl Group (e.g. CH_3-)

- In **Aldehydes**, at least **one “H”** is attached to the carbonyl (**C=O**) carbon atom.




- In **Ketones**, there are **two “C”** bonded to the carbonyl (**C=O**) carbon atom.



- The systematic names of **aldehydes** are ended by the suffix **-al** and that of **ketones** are ended by the suffix **-one**.
- They can be prepared by the controlled **oxidation of alcohols**.



Carboxylic Acids ($\text{R}-\text{COOH}$)

Carboxylic Acids contain the carboxyl group. Often written as ($-\text{COOH}$) attached to a carbon of an alkyl group (R):



Carboxylic Acids

- Carboxylic acids are weak acids, they can be produced by **oxidation of alcohols**. Under appropriate conditions, the aldehyde may be isolated as the first product of oxidation, as in the sequence




where (O) represents any **oxidizing agent** that can provide oxygen atoms.

- The systematic names for carboxylic acids always ends with ***-oic acid***

Esters (R-COOR')

➤ **Esters** are compounds in which the H-atom of a carboxylic acid is replaced by a carbon-containing group (R'):




- Esters are the products of reactions between carboxylic acids and alcohols.
- Esters are responsible for the **pleasant aroma (odor or smell)** of fruits and perfumes.




Esters

- ▶ Esters can be synthesized by **Condensation Reactions** of carboxylic acids with alcohols:




- ▶ The name of any ester consists of the name of the “R” group coming from the alcohol followed by the name of the group coming from the carboxylic acid, with the ***-ic*** replaced by ***-ate***.

For example, the ester formed from ethyl alcohol, $\text{CH}_3\text{CH}_2\text{OH}$, and **butanoic** acid, $\text{CH}_3(\text{CH}_2)_2\text{COOH}$, is:




Saponification Reaction of Esters

- The hydrolysis of an ester in the presence of a base is called **saponification**, a term that comes from the Latin word for soap, *sapon*.
 - ✓ Naturally occurring esters include fats and oils, and in making soap an animal fat or a vegetable oil is boiled with a strong base.
 - ✓ The resultant soap consists of a mixture of salts of long-chain carboxylic acids (called fatty acids), which form during the saponification reaction.



Amines ($\text{R}-\text{NH}_2$) and Amides ($\text{R}-\text{CO}-\text{NH}_2$)



Amine




Amide


- **Amines** are compounds in which one or more hydrogen atoms of ammonia (NH_3) are replaced by alkyl groups (R):



Ethylamine



Trimethylamine




Phenylamine
Aniline


- Amines are the most common **Organic Bases**.

Amines & Amides

➤ **Amides** are compounds which contain a carbonyl group (**C=O**) attached to “**N**” atom:




Ethylamide




Phenylmethanamide
Benzamide

➤ An amines with at least one “**H**” bonded to “**N**” undergo a condensation reaction with a carboxylic acid to form amide:



➤ **The amide linkage:** is the functional group in proteins, it links amino acids together to form “**polypeptides**”.















Common Functional Groups



Functional Group	Compound Type	Suffix	Structural Formula	Example
				Systematic Name (common name)
	Alkene	-ene		Ethene (Ethylene)
	Alkyne	-yne		Ethyne (Acetylene)
	Alcohol	-ol		Methanol (Methyl alcohol)
	Ether	ether		Dimethyl ether
	Aldehyde	-al		Ethanal (Acetaldehyde)
	Ketone	-one		Propanone (Acetone)
	Carboxylic acid	-oic acid		Ethanoic acid (Acetic acid)
	Ester	-oate		Methyl ethanoate (Methyl acetate)
	Amine	-amine		Ethylamine
	Amide	-amide		Ethanamide (Acetamide)

Assessment

1. Identify the type (family) of the following organic compounds



2. Identify the functional groups in each of the following compounds:



3. _____ is formed by the reaction of a carboxylic acid with an alcohol

A) aldehyde. B) ester. C) ether. D) ketone.

4. _____ is the hydrolysis of an ester using a base.

A) Saponification B) Decarboxylation
C) Detoxification D) Alcoholysis

5. Oxidation of an aldehyde produces a _____.

A) carboxylic acid. B) alcohol.
C) ester. D) ketone.

6. The common functional group in aldehydes and ketones is _____.

A) hydroxyl group. B) phenol group.
C) ether group. D) carbonyl group.

7. The Product of the reaction between a carboxylic acid and an amine is _____.

A) aldehyde. B) amide. C) ester. D) ketone.

8. _____ are used as medical anesthetics.

9. _____ are responsible for the pleasant aroma of fruits.

10. The suffix _____ is used at the end of esters names.

11. The controlled oxidation of alcohols produces _____, while further oxidation produces _____.

12. _____ are the most common organic bases.